

Risk factors and prevalence of periodontitis in community-dwelling elders in Mexico

S. Aída Borges-Yáñez¹, María Esther Irigoyen-Camacho² and Gerardo Maupomé³

¹Facultad de Odontología, Universidad Nacional Autónoma de México, México D.F., México; ²Departamento de Atención a la Salud, Universidad Autónoma Metropolitana Xochimilco, México D.F., México; ³Oral Health Research Institute, Indiana University School of Dentistry, Indianapolis, IN, USA

Borges-Yáñez SA, Irigoyen-Camacho ME, Maupomé G. Risk factors and prevalence of periodontitis in community-dwelling elders in Mexico. *J Clin Periodontol* 2006; 33: 184–194. 10.1111/j.1600-051X.2006.00897.x. © Blackwell Munksgaard, 2006.

Abstract

Aim: The objective of this study was to determine whether an association existed between chronic systemic diseases/conditions, risk factors common in old age, and the extent and severity of chronic periodontal disease.

Materials and Methods: Sociodemographic and lifestyle characteristics were examined by contrasting rural, urban-marginal, and urban social environments in Central Mexico. Data were analysed with Analysis of Variance, χ^2 tests, and multivariable logistic regression.

Results: A total of 473 adults 60 years old and over were interviewed; 315 were also examined and underwent laboratory assays (participation rate, 66%); women, 62%; mean age 73 ± 8 years; 23% edentulous. The distribution of periodontitis by sociodemographic variables showed differences across locales (73% low-urban, 57% middle-urban, 29% rural). The regression model indicated that periodontitis was more frequently associated with low-urban locale, higher systolic blood pressure, higher body mass index, and worse calculus readings, with an interaction whereby being obese and having a high calculus index was associated with a high probability of having periodontitis.

Conclusions: Overall periodontal conditions were fair. While we identified oral, systemic, and social variables that modulated the experience of periodontitis, it would appear that urban, low social class elders appeared to have worse periodontal conditions.

Key words: calculus; cardiovascular disease; diabetes; elderly; hypertension; Mexico; periodontitis; systemic disease.

Accepted for publication 29 December 2005

Moderate and severe periodontitis are defined variously, using different measurement methods and operational definitions. Despite this inconsistency in definition, the prevalence of severe periodontitis is known to be relatively high in the open population – 11–15% in adult populations in both developed (Albandar et al. 1999) and developing countries (Jiménez Garcia et al. 1995, Gjermo et al. 2002, Dowsett et al. 2001, Ronderos et al. 2001, Borges-Yáñez et al. 2004, Taani 2004). Reports from several countries have shown that the prevalence of periodontitis in older adults, defined as at least one site with clinical attachment loss (CAL) > 4 mm, is moderate to high. For example, pre-

valence was 48–55% in Chinese adults (Corbet et al. 2001); 4–59% in elderly subjects in the US Pacific Northwest (Persson et al. 2004); and 68% in the National Survey of Oral Health in US Employed Adults and Seniors (Albandar 2002b). Higher figures were found in some cases, such as 100% among 50–59 year-olds in Thailand (Baelum et al. 2003) and 91% in elderly subjects (mean age 85 years) in the US Midwest (45% with one or more sites with loss of attachment (LOA) ≥ 6 mm, and 15% with CAL ≥ 8 mm) (Levy et al. 2003). In Germany (Mack et al. 2004), the prevalence of periodontal disease (at least one periodontal pocket ≥ 4 mm) was between 62% and 85% in persons

60–79 years old. Among Mexican Americans in the US, 15% of 60–69 year olds and 9.3% of persons older than 70 years had at least four sites with CAL ≥ 5 mm and at least one site with pocket depth ≥ 4 mm (Borrell et al. 2004). When periodontal status is examined according to the Community Periodontal Index of Treatment Needs (CPITN) scores, 70% of persons 60–69 years old in the United Kingdom had CAL > 3.5 mm (Morris et al. 2001). In Finland, 11% had at least one pocket ≥ 6 mm (CPITN = 4) (Ajwani et al. 2001). In Hong Kong, the percentage of institutionalized dentate subjects with shallow pockets and deep pockets (CPITN score) were 37% and 20% respectively (Lo et al. 2004). In

Jordan, shallow (19%) and deep (11%) pockets were found in the 50–60-year-old age group (Taani 2004). Severe CAL is generally confined to a minority (Locker et al. 2000).

Limited information is available from Latin American and Mexican populations (Jiménez García et al. 1995, Borges-Yañez et al. 2004). In a Brazilian study, 100% of persons 60–69 years old had at least one site ≥ 4 mm, with worse readings associated with lower socioeconomic level, smoking, diabetes, gender, and dental visits (Susin et al. 2004). A report presenting results derived from outpatients at a dental school clinic in Mexico found 83% extent and 3.2 mm severity in persons 60 years and older (Jiménez García et al. 1995). The prevalence of moderate periodontitis (at least one site > 4 mm) in all age groups was 43%.

In older populations, the relative contributions to periodontal problems of systemic and local risk factors are unclear. At the local level, risk factors such as calculus (Albandar 1990, Albandar et al. 1996) contribute to the development of periodontal diseases in adults. Subgingival bacteria associated with periodontal/gingival conditions have been found in various countries (Haffajee et al. 2004). Even though systemic/general risk factors for periodontitis are not markedly different across young adult and elderly populations, certain factors affect the severity and distribution of periodontitis and undermine treatment outcomes in older populations. These include lifestyle characteristics, poor attitudes toward dental health, low socioeconomic and educational levels (Dolan et al. 1997, Norderyd & Hugoson 1998, Elter et al. 1999, Paulander et al. 2003), restricted access to dental services, rural residence (Craig et al. 2001, Sbaraglia et al. 2002), presence of systemic diseases (Noack et al. 2000, Buhlin et al. 2003, Persson et al. 2003, Willershausen et al. 2003, Khader et al. 2004, Yoshihara et al. 2004), acute medical conditions such as infectious respiratory diseases (Scannapieco et al. 2003), polypharmacy, poor nutritional status, obesity (Saito et al. 2001, Wood et al. 2003) mental disorders (Teng et al. 2003), and smoking (Bennet & Reade 1982, Genco 1996). More disconcerting for populations at high risk of cardiovascular complications, such as the elderly, is the fact that (Lopez et al. 2004) several studies have reported epidemiological associations between perio-

odontitis and cardiovascular disease (Mattila et al. 1989, DeStefano et al. 1993, Beck et al. 1996, Buhlin et al. 2002, Willershausen et al. 2003). The reason for such associations is unknown; the relationship might be weak (Hujoel 2002), because periodontitis and cardiovascular disease (Armitage 2000) share common risk factors (such as smoking and diabetes) (Beck et al. 1998, Hujoel et al. 2002) that can confound the association (Katz et al. 2002).

In Mexico, the National Survey of Chronic Diseases (Secretaría de Salud 1995) showed that the most frequent diseases among persons 60 years old and older were hypertension, diabetes mellitus, and cardiovascular conditions. The highest prevalence of diabetes was found in persons 70–79 years old (22.4%) (Olaiz et al. 2003) and also was high in men (20%) and women (25%) 60–69 years old. Twenty-four per cent of the Mexican population had hypertension in 1993 (Secretaría de Salud 1995), rising to 30.7% (Olaiz et al. 2003) in 2000. Furthermore, the Mexican population has a genetic predisposition to metabolic syndrome, diabetes mellitus type 2, and some dyslipidemias (Stern et al. 1991), with a high prevalence of the latter (Gonzalez et al. 1999, Aguilar-Salinas et al. 2001). The high-fat, high-carbohydrate diet, considerable alcohol intake, frequent tobacco consumption, and sedentary lifestyle in an increasingly larger proportion of the Mexican population have been identified as environmental risk factors for some chronic diseases (Munoz et al. 2000). While the availability of epidemiologic information on periodontal conditions is sparse in Mexico, knowledge of systemic/general risk factors contributing to the overall experience of periodontal/gingival conditions is even more limited. The objective of the present study was to determine whether an association existed between some systemic/general risk factors (as defined through biochemical indicators derived from laboratory tests and/or medical history), mouth-level risk factors for periodontitis, and the prevalence of moderate to severe periodontitis in persons 60 years old and older in Central Mexico.

Materials and Methods

The study was performed in accordance with the Helsinki Declaration of Human Studies, and the local Research and

Ethics Committee authorized the protocol of this study.

This cross-sectional study included study participants from three different populations (urban areas of middle and low income level, and a rural area) included in the CRONOS Study (Cross-Cultural Research on Nutrition in the Older Adult Study Group), funded by the European Economic Community. The main objective of CRONOS was to describe the nutritional situation of older people in urban and rural communities in developing countries, from transcultural and multicentric perspectives (Van Staveren 1988).

Setting

Middle-urban group

The Centro Urbano *Presidente Miguel Alemán* is a large housing development in a middle-class neighbourhood in Mexico City. The elderly residents are retired civil servants who live in small apartments bought from the federal government during the 1960s. This urban group is classified as middle social class.

Low-urban group

The *Isidro Fabela* neighborhood is on the southern outskirts of Mexico City. This peri-urban community began as an illegal settlement of squatters who migrated to the city over many years. Such rural migrants settled irregularly in an area with no public services or well-defined social organization. Over the years, public services were introduced and the community was annexed to the city in bits and pieces. The residents' characteristics of schooling, employment, and social structure define this group as low urban social class.

Rural group

The *Valley of Solis* is a rural region with several villages, located in the Northwest area of the State of Mexico in the central region of the country, 200 km from Mexico City. Agriculture (corn and wheat) is the main economic activity. Public services are limited; the maximum level of schooling attained and the social structure prevalent in the area define this group as low rural social class.

Sample design

Censuses were performed in the three communities to identify all persons 60

years old and over, and three sample frameworks were designed by systematic random selection. The total numbers of subjects 60 years and older in each community were middle-urban 464, low-urban 332, and rural 469. Sample sizes were 139 (30%), 179 (54%), and 155 (33%), respectively. Only the persons living permanently within the communities were interviewed; all were interviewed in Spanish. All participants were living independently, and no respondent was unable to respond verbally to the interview.

Data collection

The design, purpose, and activities of the study were thoroughly explained in person by staff in the first visit. Informed consent was obtained to collect data using a structured interview, a dental exam, and laboratory assays. Pilot studies were carried out in 20 persons (not included in the final study sample) to assess acceptability and clarity of the questionnaire components and to determine the time it would take to complete the interview with the instrument and the dental exam under field conditions. A previously standardized dentist performed the oral examinations with a flat dental mirror and a dental probe, with the patient sitting on a chair under natural light. Also, examiners were calibrated during the study, except for the recorders. To reduce systematic and random measurements errors, 10% subjects were recorded twice. No more than five subjects were examined daily by each examiner to avoid fatigue. Several oral health status variables were assessed, but for the present analyses only periodontal disease data will be reported in detail.

This report includes the results from 473 participants, of whom 315 agreed to a dental examination and to give a blood sample for laboratory tests. (In a separate analysis, no significant differences were observed in the sociodemographic characteristics in the group with and without biochemical analysis (data not reported). The participants received written information about the project, provided informed consent, and were examined at home. The participants received written results from the lab tests, and clinical advice was given when appropriate.

A structured interview was conducted in the participant's home. The question-

naire used in the interview assessed demographic and socioeconomic information, family health history, personal medical history, and lifestyle factors such as smoking and physical activity. The structured interview was carried out by a trained interviewer, collecting data on gender, schooling (highest grade attained in formal education), current smoking status (whether the subject stated that s/he smoked every day; number of cigarettes smoked every day), former smoker status (answer to the question: *Have you smoked in the past?*), marital status (single/ married/ widowed/ divorced), and other data. Dietitians performed anthropometric measurements and conducted dietary histories and nutritional assessments (Nelson et al. 1989). Participants' history of myocardial infarction (MI), cerebrovascular disease (CVD), arthritis or arthrosis, osteoporosis, diabetes mellitus (Type 2), and hypertension was established by direct inquiry, by clinical examination, and/or through laboratory assays.

Laboratory tests

At a primary care facility close to their homes, participants had blood specimens taken once after a 12–14 h fast for determination of plasma lipids, lipoproteins, insulin, and glucose concentrations.

Plasma glucose was analysed by the glucose oxidase method (Boehringer Mannheim, Germany). The plasma concentrations of total cholesterol and triglycerides were determined by enzymatic methods (Boehringer). High-density lipoprotein (HDL) cholesterol was measured after precipitation of very low-density lipoproteins and low-density lipoprotein (LDL) by the phosphotungstate method (Boehringer). LDL cholesterol was estimated by the Friedewald formula. The laboratory performing the tests (Instituto Nacional de Ciencias Médicas y de la Nutrición "Salvador Zubirán," Ministry of Health, México) participates in the Lipid Standardization Program of the American College of Pathology.

Definition of variables

Normal glucose tolerance (NGT) was defined as fasting blood glucose < 115 mg/dl and no history of diabetes reported on questionnaire. Diabetes was defined as a previous diagnosis of dia-

betes by World Health Organization (World Health Organization 1985) criteria or a fasting blood glucose value > 140 mg/dl with no previous history of diabetes.

Systolic (1st phase) and diastolic (5th phase) blood pressures were measured to the nearest even digit with an electric sphygmomanometer with the subject in the supine position after a 5-min. rest. Hypertension was diagnosed if blood pressure readings were greater than 160/90 mm Hg (Bots et al. 1991).

The anthropometric measurements were performed by a trained nutritionist between 8 and 11 AM. Participants removed their shoes and upper garments and donned an examining gown. Height was measured to the nearest 0.5 cm. Body weight was measured on a balance (calibrated daily) and recorded to the nearest 0.1 kg. Body mass index (BMI) was calculated as weight (kg) divided by height squared (m²) and was used as an index of overall adiposity. The cutoff point was 30 or more.

An assortment of dental variables was collected. Periodontitis was measured in terms of the LOA; subjects classified with moderate or severe disease had at least two sites with LOA 4 mm or more (Albandar et al. 1999). Severe periodontitis was considered when at least one site had LOA ≥ 6 mm. Periodontal assessments were performed in the mid-buccal and mesio-buccal sites of all teeth present in contra-lateral quadrants. (Quadrants were randomly selected, or quadrants with more teeth present were selected if substantial loss was apparent.) Attachment levels were measured using a Michigan probe (Hu-Friedy, Chicago, IL, USA). Third molars were not examined. The utilization of dental services in the last 12 months (Yes/No) did not distinguish between various purposes of dental visits but rather identified dichotomic use – non-use of services.

Statistical analyses

Statistical analyses were performed using the NCSS Package[®] (Kaysville, UT, USA). First, bivariate tests were done to describe the general features of the study populations. One-way analysis of variance was used to establish whether an association existed between (i) systolic blood pressure, glucose, cholesterol, CHDL, LDL, hypertriglyceridemia, and missing teeth, and (ii) presence or not of moderate and severe perio-

dontitis. χ^2 tests were used for (i) tobacco consumption; residence in a middle-urban, low-urban, or rural community; gender; dental plaque; and dental calculus, and (ii) presence or not of moderate and severe periodontitis.

For analysis purposes (logistic model), the dependent variable (periodontitis) was dichotomized. The multivariable model (logistic regression) was designed to identify the joint effect of local and systemic/general variables that could have modified the presence of severe or moderate periodontitis. We used the effect likelihood ratio test (LR) to ascertain statistical significance of each variable. Interactions were explored.

Results

Descriptive results

Table 1 shows population characteristics. Men made up between 30% and 46% of the three communities. Mean age was 71 years in the low-urban community and 73 in the middle-urban community and the rural community ($F = 3.2, p = 0.04$). Significant differences in the distribution by age and gender were found across the three locales ($p < 0.05$). The marital status distributions were different among the three locales, more widows and single/divorced persons live in the middle urban locale. Differences were observed in level of schooling across the three

communities, and in level of schooling for each gender between the rural community and the middle-urban community. The largest proportions of men having university studies (32%) and women having completed high school (30%) were found in the middle-urban community. The rural community had the largest proportions of illiterate men (36.7%) and women (63.1%). The use of dental services was also different across locales: in the rural group we found the lowest utilization (13.4%), while the urban group had the highest (44.1%) (Table 1). The prevalence of moderate or severe periodontitis was 50.7%.

While demographic information is included for the whole sample (473),

Table 1. Distribution of socio-demographic characteristics by locale of residence

	Locale						Total		χ^2	p
	Middle-urban		Low-urban		Rural		N	%		
	N	%	N	%	N	%				
Gender										
Male	54	30.2	52	37.4	71	45.8	177	37.4	8.6	0.013
Female	125	69.8	87	62.6	84	54.2	296	62.6		
Total	179	100.0	139	100.0	155	100.0	473	100.0		
Marital status									67.1	<0.001
Married	56	31.8	84	61.8	111	72.5	251	54.0		
Widowed	83	47.2	44	32.4	36	23.5	163	35.1		
Single/divorced	37	21.0	8	5.9	6	3.9	51	11.0		
Total	176	100.0	136	100.0	153	100.0	465	100.0		
Schooling									281.1	<0.001
Illiterate	6	3.4	56	40.6	80	52.3	142	30.2		
Elementary incomplete	29	16.2	47	34.1	69	45.1	145	30.9		
Elementary complete	30	16.8	29	21.0	4	2.6	63	13.4		
Middle and higher	114	63.7	6	4.3	0	.0	120	25.5		
Total	179	100.0	138	100.0	153	100.0	470	100.0		
Smoking									6.7	0.03
No	144	81.8	123	89.1	140	90.9	407	87.0		
Yes	32	18.2	15	10.9	14	9.1	61	13.0		
Total	176	100.0	138	100.0	154	100.0	468	100.0		
Past smoking									3.1	0.52
No	96	54.5	82	59.4	90	59.2	268	57.5		
Yes, cigarette	80	45.5	56	40.6	61	40.1	197	42.3		
Yes	0	.0	0	.0	1	.7	1	0.2		
Total	176	100.0	138	100.0	152	100.0	466	100.0		
Age									9.2	0.01
60-79	141	78.8	119	85.6	110	71.0	370	78.2		
80+	38	21.2	20	14.4	45	29.0	103	21.8		
Total	179	100.0	139	100.0	155	100.0	473	100.0		
Diabetes									11.9	0.02
No	114	77.6	61	62.9	72	80.9	247	74.2		
Yes	28	19.0	33	34.0	17	19.1	78	23.4		
Undetermined	5	3.4	3	3.1	0	.0	8	2.4		
Total	147	100.0	97	100.0	89	100.0	333	100.0		
Use of dental services in the last 12 months									30.01	<0.001
No	62	55.9	63	67.7	123	86.6	248	71.7		
Yes	49	44.1	30	32.3	19	13.4	98	28.3		
Total	111	100	93	100	142	100	346	100		
Moderate or severe periodontitis									48.5	<0.001
No	52	43	27	26.7	101	70.6	180	49.3		
Yes	69	57	74	73.3	42	29.4	185	50.7		
Total	121	100	101	100	143	100	365	100		

the periodontal examination considered only 365 (77%) persons 60 years old and over with at least one tooth present; for the logistic regression analysis, 170 persons who had complete records were included.

We compared sociodemographic variables, BMI, and periodontitis between subjects included and not included in the logistic regression analysis. No statistically significant differences were found for mean age, gender, schooling, marital status, BMI, smoking, or distribution of periodontitis; only differences in the distribution by locale were found ($p = 0.012$). No differences were found for blood pressure, cholesterol, glucose, LDL, and HDL between edentulous and non-edentulous. Overall, 22.8% of participants were edentulous and were excluded from further analyses in the present report.

Of note are the facts that only 10% of the 365 participants were current smokers; no differences were found by locale.

The highest proportion with previous diagnosis of diabetes or a fasting blood glucose value > 140 mg/dl was observed in the low-urban community (35.2%) and the lowest in the rural community (17.9%) ($\chi^2 = 6.1, p = 0.045$).

Mean glucose levels were higher in the low-urban group (119.1 mg/dl), followed by the rural group (92.9 mg/dl) and the middle-urban group (80.2 mg/dl) ($F = 10.2, p < 0.001$). Mean cholesterol value was 213.8 mg/dl, with the highest value in the middle-urban group (225 mg/dl) and the lowest in the rural group (200.5 mg/dl) ($F = 8.1, p < 0.001$). HDL mean values were similar in the rural and the middle-urban group (48 mg/

dl), and lower in the low-urban group (42.9 mg/dl) ($F = 5.1, p = 0.007$). LDL mean values were higher in the middle-urban group (157.1), followed by the low-urban group (144.3) and the rural group (130.4) ($F = 10.3, p < 0.001$). The group showing the highest mean BMI score was the low-urban (27.3), and the lowest was the rural group (23.2) ($F = 23.7, p < 0.001$). Hypertension higher than 160/90 mm Hg was more prevalent in the low-urban community (46%), followed by the middle-urban community (31%) and the rural community (27.9%) ($\chi^2 = 9.7, p = 0.008$). The highest systolic blood pressure values were observed in the middle-urban community (138.3 mm Hg) and the lowest in the rural community (120.3 mm Hg), while diastolic blood pressure was similar among the three communities (81.7 mm Hg) ($F = 2.6, p = 0.09$).

The distribution of periodontitis by sociodemographic variables showed that differences were significant across locales (73% periodontitis in the low-urban group, 57% in the middle-urban group, and 29% in the rural group ($\chi^2 = 48.5, p < 0.001$)).

In general, we did not observe differences in terms of periodontitis across the various levels of utilization of dental services in the prior 12 months. For example, of those that had not used dental services, 49.4% had moderate or severe periodontitis, compared to 55.4% who had used dental services ($\chi^2 = 0.97, p = 0.324$). Differences across locales were not significant.

Regarding the mean values for biochemical markers, BMI, age, number of teeth, plaque, and calculus across the

presence or absence of moderate and severe periodontitis, we found that mean age (73.2 years) and lower number of teeth (14.6), and higher values for systolic blood pressure (132.8 mm Hg), plaque (1.2), and calculus (1.4) were significantly higher in the severe or moderate periodontitis group than in the no periodontitis group (Table 2). The distribution of periodontitis showed marginally significant differences ($\chi^2 = 2.8, p = 0.09$) in the diabetic group (61.5%) versus the non-diabetic group (49.5%). Current smoking was not associated with periodontitis ($\chi^2 = 2.2, p = 0.13$).

Multivariate model results

Our multiple logistic regression model used moderate and severe periodontal disease as the outcome variable (Table 3). The following variables were significant: locale (likelihood ratio (LR) $p = 0.0031$), systolic blood pressure (LR $p = 0.017$), and calculus (LR $p = 0.0625$). An interaction between BMI and calculus was detected (LR $p = 0.05$). To explore the interaction, both variables were dichotomized: the cutoff point for BMI was ≥ 30 and the model used the mean value for calculus. In this way, four groups were formed: high BMI and high calculus; high BMI and low calculus; low BMI and high calculus; and low BMI and low calculus. The probability estimated by the model to have periodontitis was 0.68, 0.34, 0.60 and 0.43, respectively, suggesting that being obese and having a high calculus index was associated with a high probability of having periodontitis.

Table 2. Mean values for laboratory assay markers, body mass index, age, number of teeth, dental plaque, and dental calculus across presence of periodontitis

	No periodontitis			Moderate or severe periodontitis			Total			<i>t</i>	<i>p</i>
	<i>N</i>	Mean	SD	<i>N</i>	Mean	SD	<i>N</i>	Mean	SD		
Age	180	71.5	8.26	185	73.20	8.3	365	72.35	8.30	3.9	0.048
BMI	152	25.6	4.84	169	25.73	4.9	321	25.67	4.87	0.06	0.814
Teeth	180	18.5	8.27	185	14.62	7.6	365	16.55	8.16	22.2	<0.001
Systolic BP (mm Hg)	144	126.8	21.06	137	132.88	24.2	281	129.78	22.81	5.0	0.026
Diastolic BP (mm Hg)	142	81.4	19.12	134	82.07	18.2	276	81.71	18.63	0.09	0.758
Glucose (mg/dL)	117	93.1	45.46	126	102.86	48.4	243	98.17	47.15	2.6	0.108
Cholesterol (mg/dL)	117	214.6	40.42	126	213.04	42.5	243	213.79	41.43	0.09	0.769
Triglycerides (mg/dL)	117	155.5	96.02	126	166.18	125.7	243	161.04	112.29	.55	0.460
HDL (mg/dL)	117	48.1	12.49	126	46.15	12.7	243	47.1	12.61	1.5	0.227
LDL (mg/dL)	115	144.9	40.16	124	144.54	40.1	239	144.76	40.04	0.01	0.930
Dental calculus	168	.83	0.65	171	1.21	0.91	339	1.02	.814	19.6	<0.001
Dental plaque	168	1.0	0.61	171	1.38	0.86	339	1.20	0.77	20.1	<.001

BMI, body mass index; BP, blood pressure; HDL, high-density lipoprotein; LDL, low-density lipoprotein.

Table 3. Logistic regression model for periodontitis (dependent variable: moderate or severe periodontal disease, two or more sites with LOA ≥ 4 mm)

Term	Estimate	Standard error	Probability $> \chi^2$	OR	CI lower 95%	CI upper 95%
Intercept	-4.3149	2.4095	0.0733			
Referent rural locale						
Urban middle locale	0.6416	0.4988	0.1983	1.8994	0.7164	5.1176
Urban low locale	1.8142	0.5683	0.0014	6.1361	19.4853	2.0746
Age	0.0479	0.0275	0.0813	1.0490	0.9948	1.1086
Referent male						
Gender (female)	-0.7733	0.4215	0.0665	1.1367	1.0000	1.3058
Systolic blood pressure	0.0204	0.0088	0.0199	1.0206	1.0036	1.0389
BMI	-0.0698	0.0442	0.1142	0.9325	0.8524	1.0170
Referent no diabetes M						
Diabetes	0.0764	0.4553	0.8668	1.0794	0.4390	2.6473
Number of teeth present	0.0447	0.0249	0.0725	1.0457	0.9965	1.0991
Calculus (BMI) [*] (calculus)	0.2319	0.2507	0.3548	1.2610	0.7779	2.0964
Referent non-smoker						
Current smoker	0.5806	0.6537	0.3744	1.7872	0.4974	6.6898
Referent no dental visit last 12 months						
Dental visit last 12 months	-0.2221	0.4204	0.5974	0.8009	0.3490	1.8296

Pseudo- $R^2 = 0.2097$; $\chi^2 = 48.17$; $p < 0.0001$.

LOA, loss of attachment.

Discussion

Periodontal conditions

Periodontal measurements

Of the elders examined, 28% experienced moderate periodontitis (at least two sites with LOA ≥ 4 mm). Comparing this prevalence with previously published reports is difficult due to the diversity of disease definitions, classification categories, and cutoff points that appear in the literature (Kingman & Albandar 2002). In the absence of surveys in Mexican elders using the definition we used, we will contrast our findings with international studies that used LOA to estimate periodontitis prevalence. In Germany, the prevalence of periodontal disease (presence of at least one periodontal pocket ≥ 4 mm) in 60–69-year-old persons was 85% in men and 71% in women (Mack et al. 2004), which was higher than the prevalence we observed. A study in Thailand detected 100% of 50–59 year olds having at least one site with LOA ≥ 4 mm (Baelum et al. 2003), while a study in Brazil found that 100% of adults aged 60–69 years old presented with LOA ≥ 4 mm (Susin et al. 2004). Allowing for some lassitude in terms of methods being somewhat different, Argentinian (34%, moderate and 39%, severe periodontitis) and Brazilian data (49%, moderate and 18%, severe perio-

odontitis) offer similar prevalence levels to our results (Gjermeo et al. 2002).

Studies in American populations also showed a higher prevalence of LOA than our study. For example, Albandar (2002b) found higher prevalence in the National Survey of Oral Health in US Employed Adults and Seniors, 1985–1986, with 51.7% of seniors having attachment loss ≥ 6 mm. Levy et al. (2003) found that 91% of the subjects (mean age 85 years) had at least one site with LOA ≥ 4 mm, and 45% had one or more sites with LOA ≥ 6 mm. In a study of Mexican-Americans aged 50 years and older, the prevalence of periodontitis was 9.4%, defining periodontitis as at least four sites with LOA ≥ 5 and at least one site with pocket depth ≥ 4 mm (Borrell et al. 2004). Comparing the prevalence with previously published reports is difficult due to the diversity of disease definitions, classification categories, and cutoff points that appear in the literature, (Kingman & Albandar 2002), in the comparison above described the same cut-off point is used (LOA ≥ 4). However, there are differences in the number of sites and teeth examined among studies and age groups. Borrell & Papapanou (2005) addressed that there is a need to standardize the ‘‘case’’ definition of periodontitis, so more valid comparisons among different populations and analysis of risk factors can be done.

Calculus

Plaque and calculus are established risk factors for periodontitis (Abdellatif & Burt 1987, Page et al. 2002), and we found them to be associated with periodontal disease. The low urban group had the highest prevalence of periodontal disease and also showed more plaque and calculus than the other two groups. However, no difference in plaque and calculus accumulation was found between the middle-urban and rural groups. Furthermore, an interaction was detected with BMI (see below).

Systemic/general health conditions

Cardiovascular conditions

Several cardiovascular risk factors were present in a large proportion of the elders. About a quarter of them presented high levels of triglycerides and cholesterol, and close to one-fifth were obese. Our findings are similar to those of the National Health Survey in Mexico (Velázquez et al. 2002), where 21.8% of elders (70–79 years old) were obese. A large Mexican survey found that adults in 417 cities had a poor lipid profile (HDL cholesterol was low in about a third of the population) and a high prevalence of insulin resistance (Aguilar-Salinas et al. 2001). In recent years, there has been an increase in the prevalence of cardiovascular risk factors and diabetes in the Mexican population overall (Lara-Rodriguez et al. 1996; Velázquez et al. 2002). The reasons for this deterioration are complex and relate to environmental factors, such as lifestyle (predominantly poor patterns of physical activity and diet), as well as genetic factors (Escobedo-de la Pena & Santos-Burgoa 1995, Flores-Martinez et al. 2004).

Blood pressure

High blood pressure was highly prevalent in the population studied – 61.2% were hypertensive. This prevalence was higher than the values obtained in the National Chronic Disease Survey (1993) (Tapia-Conyer 1993) and the National Health Survey in Mexico (2000), where 51.9% of the 70–79 year olds had hypertension (Olaiz et al. 2003). An additional problem in the Mexican population is that only 20% of patients had their hypertension situation adequately managed (Velázquez et al. 2002). We found a positive association

between systolic blood pressure and periodontal disease, but not with diastolic blood pressure, as did a study in Italian patients (Angeli et al. 2003). Left ventricular hypertrophy, which is a frequent finding in hypertensive patients (Cuspidi et al. 2005), was an independent predictor of periodontal disease. But when this variable was incorporated in the model, systolic blood pressure was no longer significant (Angeli et al. 2003).

Some epidemiological studies have shown a relationship between periodontal disease or tooth loss and high blood pressure (Taguchi et al. 2004). Data from the Framingham Heart Study showed that the relationship of blood pressure and risk of coronary heart disease changes with age. In younger subjects, diastolic blood pressure is a stronger predictor of coronary heart disease; however, in older people, diastolic and systolic pressure, and pulse were comparable risk factors of coronary heart disease (Franklin et al. 2001). It appears possible that the relationships of diastolic and systolic blood pressure and periodontitis change with age.

The reason for the association between hypertension and periodontitis is not clear. It may be that development of microcirculation dysfunction derived from hypertension affects periodontal tissues, perhaps coupled with arteriolar and capillarity rarefaction that can induce ischemia (Chapple et al. 2000, Serne et al. 2001). In hypertensive patients, changes in the vessel layers had been detected as a thickening of the intima and elastic layer, and a decrease in the lumen of vessels irrigating the periodontal membrane (Castelli et al. 1978). With the homeostasis of the periodontal tissues altered, the inflammatory conditions present in periodontitis could increase damage to periodontal tissues. Alternatively, the association of high blood pressure with periodontal disease may be related to side effects of drugs used to treat high blood pressure, since the β -adrenoceptor antagonists tend to produce xerostomia (Nederfors 1996). The OH:SALSA aging study detected a 20% reduction in salivary output associated with drugs used for high blood pressure (Dodds et al. 2000). While xerostomia can affect oral mucosa, a relationship between xerostomia and periodontal disease has not been unequivocally established (Janket et al. 2003).

Finally, data from the NHANES III showed that C-reactive protein (CRP)

and blood pressure were positively related (King et al. 2004). The complex scenario depicting the relationship between periodontal disease and CVD risk factors is compounded by the likelihood of it being bidirectional, with a large array of immunological, inflammatory, and genetic factors intervening (Slade et al. 2003; Pussinen et al. 2004).

Diabetes

The prevalence of diabetes in 60–79 year olds was about 22% in the National Health Survey 2000, close to the 25% found in the present study. The association between LOA and diabetes mellitus was marginally significant in the bivariate analysis and not significant in the multivariate model. The relationship between diabetes and periodontitis has been found in population-based surveys (Skrepcinski & Niendorff 2000, Tsai et al. 2002) but might be less consistent in elderly persons, perhaps due to accumulated/differential tooth loss and presence of co-morbidities (Zielinski et al. 2002, Persson et al. 2003), or it might be altogether absent (Tomita et al. 2002, Persson et al. 2003). Establishing the relationship between these two chronic conditions is of particular importance considering the high prevalence of diabetes in Mexico. Questions about sharing common genetic markers, effect of periodontal treatment on glycemic control, impact of lack of good glucose control on periodontitis, and the web of interactions between these aspects and psychosocial variables in the elderly require further research.

Body mass index

The association of BMI and periodontal disease is equivocal (Al Zahrani et al. 2003, Wood et al. 2003, Torrungruang et al. 2005). NHANES III data showed an association between BMI in young subjects (18–34 year olds) and periodontal disease; however, no association was detected in the older age groups (Al Zahrani et al. 2003). A cross-sectional study in Japanese subjects showed that in the highest categories of BMI, obesity was a significant risk factor for periodontitis after adjusting for several known periodontitis risk factors (Saito et al. 2001). A case control study of periodontitis and risk markers for atherosclerosis found an association with BMI, which was higher among cases

with severe periodontal disease (Buhlin et al. 2003).

This is the first study, to our knowledge, to find an interaction between BMI and calculus. The logistic regression model showed that the group with high BMI (BMI ≥ 30) and abundant calculus had a high probability of having moderate or severe periodontitis. Longitudinal studies had showed that good oral hygiene is a highly specific predictor of periodontal stability (Lang et al. 1990). The relatively weak interaction between BMI and calculus suggests that more studies are necessary to investigate in greater detail the complex relationships between periodontal features, and local and systemic risk factors for periodontitis in the elderly. We could speculate that the presence of calculus may be a marker for use of dental services (possible surrogates for poorly defined concepts such as access to dental services, or dental IQ), accepting that the relationship could be modified by other factors.

Smoking

The proportions of current smokers and never smokers found in the present study are consistent with data from the National Health Survey 2000 in Mexico. In the present study, 10.2% were current smokers and 58.4% never smoked, corresponding to the National Survey proportions of 11.5% and 59.8%. At the bivariate analysis level we found an association between periodontitis and cigarettes smoked, resembling observations derived from studies in adults living in developed and developing countries that support a dose–response relationship (Do et al. 2003, Nishida et al. 2005). In the moderate SES group, 13.0% of elders smoked, leading to an association between tobacco use and moderate or severe periodontitis in which 86.7% of smokers were affected. In the two low socioeconomic level locales (rural or urban), overall tobacco use and mean cigarette consumption were lower to begin with – we speculate that the two features contributed to the observed lack of association in the regression model. While we acknowledge that self-reported smoking may have validity problems, other studies have not been able to detect an association between periodontal disease and tobacco in the elderly (Persson et al. 2004, Thomson et al. 2004). In fact, a study in Sri Lanka showed that the

effect of oral hygiene was more important than tobacco use as an indicator of risk of periodontitis (Amarasena et al. 2002), and another study in the USA suggested that smoking was a weak predictor of periodontitis in the elderly (Persson et al. 2005). The proportion of current smokers and the cut-off point for periodontitis (LOA ≥ 4) in the latter study and ours are very similar.

Locale of residence

Important differences among the urban and rural elderly were observed for both periodontal disease and cardiovascular risk factors. The elderly from the rural community had lower prevalence of periodontal disease than the middle-urban and low-urban groups. This finding was the opposite of findings from North America, Africa, and Scandinavia, where rural inhabitants had a higher prevalence of periodontal disease (Dolan et al. 1997, Paulander et al. 2004, Varenne et al. 2004). Furthermore, elderly community-dwelling Israelis living in rural areas had a higher mean attachment loss than those living in urban areas (Adut et al. 2004).

In the present study, the rural elderly had not only better periodontal health than their urban counterparts but also a lower prevalence of cardiovascular risk factors. While it is possible that lifestyle and diet drive these results (de Chavez et al. 1993), such differences also may be related to overall health status.

It is important to point out that very probably there are differences in the type of diet and physical activity among the three localities, since these characteristics are inherent to the geographical and socio-cultural environment of each locale. For example, the elderly living in the rural setting have more physical activity and probably a higher fibre intake than the elderly living in urban localities.

Some studies have highlighted the intimate association between periodontal disease and cardiovascular diseases, stroke, and other systemic problems (Beck et al. 1996, Geerts et al. 2004, Paquette 2004). These factors are different between the rural and the urban areas in the present study, with urban elderly experiencing higher levels of fasting glucose, prevalence of diabetes, and BMI than the rural elderly. In addition, mean cholesterol values were higher than the recommended values in the urban groups, and the rural group

had both a better lipid profile and a lower systolic blood pressure. The commonly assumed facts of sparser health services and lower life expectancy in the rural area suggest that we could be witnessing a ‘healthier survivor bias,’ but it may be that the health difference between the urban and rural areas depends on other factors – for example, psychological aspects. A study of deprivation, social class, and quality of life in British elderly showed that living in a rural area was associated with lower risk of poor morale (Breeze et al. 2005). At the present stage of our research, we are able only to speculate about other dimensions of health-related factors that could be contributing to the phenomena observed.

Socioeconomic status variables

Inequalities in health status associated with socioeconomic variables have been documented for decades in the North America, Europe, and Latin America (Feinstein 1993, Almeida-Filho et al. 2003, Comaru & Westphal 2004, Kunst et al. 2005). Higher periodontal disease prevalence has been reported in low income groups as well (Rozier & Beck 1991, Dolan et al. 1997). However, this relationship is not universal: while an analysis of social factors and periodontal diseases in the NHANES III showed that high-income Blacks had a higher prevalence of periodontitis than low-income Blacks, this type of relationship was not detected among Mexican-Americans. Conversely, lower educational attainment was associated with higher prevalence of periodontal disease in Mexican-Americans (Borrell et al. 2004). We found that the low-income urban group also had a lower level of education and showed the highest periodontal damage, compared with the middle-income urban group. Psychological aspects may play a role in these differences – Genco et al. (1999) found that variables associated with psychological stress, particularly those related with economic constraints, had a negative impact on periodontal health (Gjermeo et al. 2002), perhaps as a consequence of changes in inflammatory responses (Kiecolt-Glaser et al. 2003).

After adjusting for selected systemic and local risk factors, the effect of locale of residence remained significant. Both rural elderly and low-income urban elderly were poor, but their health status differences were considerable. It

appears that being poor in a rural area and being poor in an urban area have a different impact on cardiovascular and periodontal risk.

Conclusions

The results of the present study have good external validity considering its probabilistic sample design and the high participation rate of the elderly persons in the selected locales. Population-based studies such as this one have the advantage of reducing selection bias that is common in samples taken from hospitals or dental clinics. Conclusions from the present study, however, cannot be directly extrapolated to the entire elderly Mexican population.

In general, we found that periodontal conditions among elderly persons living in middle-urban, low-urban, and rural locales had substantial room for improvement. The poor elderly living in urban environments could have the worst of both worlds, however, through being affected by lower educational attainment and income, and the poorer diet and lifestyle of urban Mexico. The distribution of the elderly segment of the population in Mexico and its future projections (Consejo Nacional de Población 1999) indicate that the proportion of older adults will increase from 4.8% in 2000 to 6.2% in 2010 and to 13.2% by 2030. Such growth rates are similar to the trends forecasted for developed and developing countries, emphasizing the need to accurately establish the profile of oral health problems that will challenge the well-being and health care needs for this segment of the population.

Acknowledgements

This study was supported by the Consejo Nacional de Ciencia y Tecnología de México (Grant number 4323 M), and the Universidad Nacional Autónoma de México (Grant number PAPIIT IN213101).

References

- Abdellatif, H. M. & Burt, B. A. (1987) An epidemiological investigation into the relative importance of age and oral hygiene status as determinants of periodontitis. *Journal of Dental Research* **66**, 13–18.
- Adut, R., Mann, J. & Sgan-Cohen, H. D. (2004) Past and present geographic location as oral health markers among older adults. *Journal of Public Health Dentistry* **64**, 240–243.

- Aguilar-Salinas, C. A., Olaiz, G., Valles, V., Torres, J. M., Gomez Perez, F. J., Rull, J. A., Rojas, R., Franco, A. & Sepulveda, J. (2001) High prevalence of low HDL cholesterol concentrations and mixed hyperlipidemia in a Mexican nationwide survey. *Journal of Lipid Research* **42**, 1298–1307.
- Ajwani, S., Tervonen, T., Narhi, T. O. & Ainamo, A. (2001) Periodontal health status and treatment needs among the elderly. *Special Care in Dentistry* **21**, 98–103.
- Al Zahrani, M. S., Bissada, N. F. & Borawskit, E. A. (2003) Obesity and periodontal disease in young, middle-aged, and older adults. *Journal of Periodontology* **74**, 610–615.
- Albandar, J. M. (1990) Some predictors of radiographic alveolar bone height reduction over 6 years. *Journal of Periodontal Research* **25**, 186–192.
- Albandar, J. M. (2002a) Global risk factors and risk indicators for periodontal diseases. *Periodontology 2000* **29**, 177–206.
- Albandar, J. M. (2002b) Periodontal diseases in North America. *Periodontology 2000* **29**, 31–69.
- Albandar, J. M., Brown, L. J., Brunelle, J. A. & Loe, H. (1996) Gingival state and dental calculus in early-onset periodontitis. *Journal of Periodontology* **67**, 953–959.
- Albandar, J. M., Brunelle, J. A. & Kingman, A. (1999) Destructive periodontal disease in adults 30 years of age and older in the United States 1988–1994. *Journal of Periodontology* **70**, 13–29.
- Almeida-Filho, N., Kawachi, I., Filho, A. P. & Dachs, J. N. (2003) Research on health inequalities in Latin America and the Caribbean: bibliometric analysis (1971–2000) and descriptive content analysis (1971–1995). *American Journal of Public Health* **93**, 2037–2043.
- Amarasena, N., Ekanayaka, A. N., Herath, L. & Miyazaki, H. (2002) Tobacco use and oral hygiene as risk indicators for periodontitis. *Community Dentistry and Oral Epidemiology* **30**, 115–123.
- Angeli, F., Verdecchia, P., Pellegrino, C., Pellegrino, R. G., Pellegrino, G., Prosciutti, L., Giannoni, C., Cianetti, S. & Bentivoglio, M. (2003) Association between periodontal disease and left ventricle mass in essential hypertension. *Hypertension* **41**, 488–492.
- Armitage, G. C. (2000) Periodontal infections and cardiovascular disease – how strong is the association? *Oral Diseases* **6**, 335–350.
- Baelum, V., Pisuithanank, S., Teanpaisan, R., Pithpornchaiyakul, W., Pongpaisal, S., Papanou, P. N., Dahlen, G. & Fejerskov, O. (2003) Periodontal conditions among adults in Southern Thailand. *Journal of Periodontal Research* **38**, 156–163.
- Beck, J. D., Garcia, R., Heiss, G., Vokonas, P. S. & Offenbacher, S. (1996) Periodontal disease and cardiovascular disease. *Journal of Periodontology* **67**, 1123–1137.
- Beck, J. D., Offenbacher, S., Williams, R., Gibbs, P. & Garcia, R. (1998) Periodontitis: a risk factor for coronary heart disease? *Annals of Periodontology* **3**, 127–141.
- Bennet, K. R. & Reade, P. C. (1982) Salivary immunoglobulin A levels in normal subjects, tobacco smokers, and patients with minor aphthous ulceration. *Oral Surgery Oral Medicine Oral Pathology* **53**, 461–465.
- Borges-Yañez, S. A., Maupome, G., Martinez-Gonzalez, M., Cervantez-Turrubiate, L. & Gutierrez-Robledo, L. M. (2004) Dietary fiber intake and dental health status in urban-marginal, and rural communities in central Mexico. *Journal of Nutrition Health Aging* **8**, 333–339.
- Borrell, L. N., Burt, B. A., Neighbors, H. W. & Taylor, G. W. (2004) Social factors and periodontitis in an older population. *American Journal of Public Health* **94**, 748–754.
- Borrell, L. N. & Papanou, P. N. (2005) Analytical epidemiology of periodontitis. *Journal of Clinical Periodontology* **32** (Suppl. 6), 132–158.
- Bots, M. L., Grobbee, D. E. & Hofman, A. (1991) High blood pressure in the elderly. *Epidemiologic Reviews* **13**, 294–314.
- Breeze, E., Jones, D. A., Wilkinson, P., Bulpitt, C. J., Grundy, C., Latif, A. M. & Fletcher, A. E. (2005) Area deprivation, social class, and quality of life among people aged 75 years and over in Britain. *International Journal of Epidemiology* **34**, 276–283.
- Buhlin, K., Gustafsson, A., Hakansson, J. & Klinge, B. (2002) Oral health and cardiovascular disease in Sweden. *Journal of Clinical Periodontology* **29**, 254–259.
- Buhlin, K., Gustafsson, A., Pockley, A. G., Frostegard, J. & Klinge, B. (2003) Risk factors for cardiovascular disease in patients with periodontitis. *European Heart Journal* **24**, 2099–2107.
- Castelli, W. A., Diaz-Perez, R., Nasjleti, C. E. & Caffesse, R. G. (1978) Effect of renovascular hypertension on the morphology of oral blood vessels. *Oral Surgery Oral Medicine Oral Pathology* **46**, 576–582.
- Chapple, C. C., Kumar, R. K. & Hunter, N. (2000) Vascular remodelling in chronic inflammatory periodontal disease. *Journal of Oral Pathology and Medicine* **29**, 500–506.
- de Chavez, M. M., Valles, V., Blatter, F., Avila, A. & Chavez, A. (1993) [Rural and urban nutrition and its relationship to the atherogenic risk.] *Salud Publica de Mexico* **35**, 651–657.
- Comaru, F. de A. & Westphal, M. F. (2004) Housing, urban development and health in Latin America: contrasts, inequalities and challenges. *Reviews On Environmental Health* **19**, 329–345.
- Consejo Nacional de Población (1999) *El Envejecimiento Demográfico de México: Retos y Perspectivas: Por Una Sociedad Para Todas Las Edades*, 1ª edición. México: D.F. CONAPO.
- Corbet, E. F., Wong, M. C. & Lin, H. C. (2001) Periodontal conditions in adult Southern Chinese. *Journal of Dental Research* **80**, 1480–1485.
- Craig, R. G., Boylan, R., Yip, J., Bamgboye, P., Koutsoukos, J., Mijares, D., Ferrer, J., Imam, M., Socransky, S. S. & Haffajee, A. D. (2001) Prevalence and risk indicators for destructive periodontal diseases in 3 urban American minority populations. *Journal of Clinical Pharmacology* **28**, 524–535.
- Cuspidi, C., Meani, S., Fusi, V., Valerio, C., Catini, E., Sala, C., Sampieri, L., Magrini, F. & Zanchetti, A. (2005) Prevalence and correlates of left atrial enlargement in essential hypertension: role of ventricular geometry and the metabolic syndrome: the Evaluation of Target Organ Damage in Hypertension study. *Journal of Hypertension* **23**, 875–882.
- De Stefano, F., Anda, R. F., Kahn, H. S., Williamson, D. F. & Russell, C. M. (1993) Dental disease and risk of coronary heart disease and mortality. *BMJ* **306**, 688–691.
- Do, G. L., Spencer, A. J., Roberts-Thomson, K. & Ha, H. D. (2003) Smoking as a risk indicator for periodontal disease in the middle-aged Vietnamese population. *Community Dentistry and Oral Epidemiology* **31**, 437–446.
- Dodds, M. W., Yeh, C. K. & Johnson, D. A. (2000) Salivary alterations in type 2 (non-insulin-dependent) diabetes mellitus and hypertension. *Community Dentistry and Oral Epidemiology* **28**, 373–381.
- Dolan, T. A., Gilbert, G. H., Ringelberg, M. L., Legler, D. W., Antonson, D. E., Foerster, U. & Heft, M. W. (1997) Behavioral risk indicators of attachment loss in adult Floridians. *Journal of Clinical Periodontology* **24**, 223–232.
- Dowsett, S. A., Archila, L., Segreto, V. A., Eckert, G. J. & Kowolik, M. J. (2001) Periodontal disease status of an indigenous population of Guatemala, Central America. *Journal of Clinical Periodontology* **28**, 663–671.
- Elter, J. R., Beck, J. D., Slade, G. D. & Offenbacher, S. (1999) Etiologic models for incident periodontal attachment loss in older adults. *Journal of Clinical Periodontology* **26**, 113–123.
- Escobedo-de la Pena, J. & Santos-Burgoa, C. (1995) [Diabetes mellitus and the health care transition]. *Salud Publica de Mexico* **37**, 37–46.
- Feinstein, J. S. (1993) The relationship between socioeconomic status and health: a review of the literature. *Milbank Quarterly* **71**, 279–322.
- Flores-Martinez, S. E., Islas-Andrade, S., Machorro-Lazo, M. V., Revilla, M. C., Juarez, R. E., Mujica-Lopez, K. I., Moran-Moguel, M. C., Lopez-Cardona, M. G. & Sanchez-Corona, J. (2004) DNA polymorphism analysis of candidate genes for type 2 diabetes mellitus in a Mexican ethnic group. *Annals of Genetics* **47**, 339–348.
- Franklin, S. S., Larson, M. G., Khan, S. A., Wong, N. D., Leip, E. P., Kannel, W. B. & Levy, D. (2001) Does the relation of blood pressure to coronary heart disease risk change with aging? The Framingham Heart Study. *Circulation* **103**, 1245–1249.
- Geerts, S. O., Legrand, V., Charpentier, J., Albert, A. & Rompen, E. H. (2004) Further evidence of the association between perio-

- dontal conditions and coronary artery disease. *Journal of Periodontology* **75**, 1274–1280.
- Genco, R. J. (1996) Current view of risk factors for periodontal diseases. *Journal of Periodontology* **67**, S1041–1049.
- Genco, R. J., Ho, A. W., Grossi, S. G., Dunford, R. G. & Tedesco, L. A. (1999) Relationship of stress, distress and inadequate coping behaviors to periodontal disease. *Journal of Periodontology* **70**, 711–723.
- Gjermo, P., Rosing, C. K., Susin, C. & Oppermann, R. (2002) Periodontal diseases in Central and South America. *Periodontology* **2000** **29**, 70–78.
- Gonzalez, C., Stern, M. P., Gonzalez, E., Rivera, D., Simon, J., Islas, S. & Haffner, S. (1999) The Mexico City diabetes study: a population-based approach to the study of genetic and environmental interactions in the pathogenesis of obesity and diabetes. *Nutrition Reviews* **57**, S71–S76.
- Haffajee, A. D., Bogren, A., Hasturk, H., Feres, M., Lopez, N. J. & Socransky, S. S. (2004) Subgingival microbiota of chronic periodontitis subjects from different geographic locations. *Journal of Clinical Periodontology* **2000** **31**, 996–1002.
- Hujoel, P. P. (2002) Does chronic periodontitis cause coronary heart disease? A review of the literature. *Journal of the American Dental Association* **133**, 31S–36S.
- Hujoel, P. P., Drangsholt, M., Spiekerman, C. & De Rouen, T. A. (2002) Periodontitis-systemic disease associations in the presence of smoking – causal or coincidental? *Periodontology* **2000** **30**, 51–60.
- Janket, S. J., Jones, J. A., Rich, S., Meurman, J., Garcia, R. & Miller, D. (2003) Xerostomic medications and oral health: the Veterans Dental Study (part I). *Gerodontology* **20**, 41–49.
- Jiménez García, G., Herrera Echauri, R. & Borges-Yáñez, S. A. (1995) Severidad y extensión de periodontitis en sujetos adultos que acudieron a la Clínica de Admisión de la División de Estudios de Posgrado e Investigación, UNAM. *Práctica Odontológica* **16**, 29–35.
- Katz, J., Flugelman, M. Y., Goldberg, A. & Heft, M. (2002) Association between periodontal pockets and elevated cholesterol and low density lipoprotein cholesterol levels. *Journal of Periodontology* **73**, 494–500.
- Khader, Y. S., Albashairh, Z. S. & Alomari, M. A. (2004) Periodontal diseases and the risk of coronary heart and cerebrovascular diseases: a meta-analysis. *Journal of Periodontology* **75**, 1046–1053.
- Kiecolt-Glaser, J. K., Preacher, K. J., MacCallum, R. C., Atkinson, C., Malarkey, W. B. & Glaser, R. (2003) Chronic stress and age-related increases in the proinflammatory cytokine IL-6. *Proceedings of the National Academy of Sciences of the United States of America* **100**, 9090–9095.
- King, D. E., Egan, B. M., Mainous, A. G. III & Geesey, M. E. (2004) Elevation of C-reactive protein in people with prehypertension. *Journal of Clinical Hypertension (Greenwich)* **6**, 562–568.
- Kingman, A. & Albandar, J. M. (2002) Methodological aspects of epidemiological studies of periodontal diseases. *Periodontology* **2000** **29**, 11–30.
- Kunst, A. E., Bos, V., Lahelma, E., Bartley, M., Lissau, I., Regidor, E., Mielck, A., Cardano, M., Dalstra, J. A., Geurts, J. J., Helmert, U., Lennartsson, C., Ramm, J., Spadea, T., Stroenegger, W. J. & Mackenbach, J. P. (2005) Trends in socioeconomic inequalities in self-assessed health in 10 European countries. *International Journal of Epidemiology* **34**, 295–305.
- Lang, N. P., Adler, R., Joss, A. & Nyman, S. (1990) Absence of bleeding on probing. An indicator of periodontal stability. *Journal of Clinical Periodontology* **17**, 714–721.
- Lara-Rodriguez, M. A., Benitez-Martinez, M. G., Fernandez-Garate, I. H. & Zarate-Aguilar, A. (1996) [Epidemiologic aspects of the aged at the Mexican Institute of Social Security]. *Salud Publica de Mexico* **38**, 448–457.
- Levy, S. M., Warren, J. J., Chowdhury, J., De Bus, B., Watkins, C. A., Cowen, H. J., Kirchner, H. L. & Hand, J. S. (2003) The prevalence of periodontal disease measures in elderly adults, aged 79 and older. *Special Care in Dentistry* **23**, 50–57.
- Lo, E. C., Luo, Y. & Dyson, J. E. (2004) Oral health status of institutionalised elderly in Hong Kong. *Community Dental Health* **21**, 224–226.
- Locker, D., Slade, G. D. & Murray, H. (2000) Epidemiology of periodontal disease among older adults: a review. *Periodontology* **16**, 16–33.
- Lopez, N. J., Socransky, S. S., Da Silva, I., Japlit, M. R. & Haffajee, A. D. (2004) Subgingival microbiota of Chilean patients with chronic periodontitis. *Journal of Periodontology* **75**, 717–725.
- Mack, F., Mojon, P., Budtz-Jorgensen, E., Kocher, T., Splieth, C., Schwahn, C., Bernhardt, O., Gesch, D., Kordass, B., John, U. & Biffar, R. (2004) Caries and periodontal disease of the elderly in Pomerania, Germany: results of the Study of Health in Pomerania. *Gerodontology* **21**, 27–36.
- Mattila, K. J., Nieminen, M. S., Valtonen, V. V., Rasi, V. P., Kesaniemi, Y. A., Syrjala, S. L., Jungell, P. S., Isolouma, M., Hietaniemi, K. & Jokinen, M. J. (1989) Association between dental health and acute myocardial infarction. *BMJ* **298**, 779–781.
- Morris, A. J., Steele, J. & White, D. A. (2001) The oral cleanliness and periodontal health of UK adults in 1998. *British Dental Journal* **191**, 186–192.
- Munoz, D. C., Chavez, V. A., Chavez, M. M. & Eichen, V. I. (2000) Sale of street food in Latin America. The Mexican case: joy or jeopardy? *World Review of Nutrition and Dietetics* **86**, 138–154.
- Nederfors, T. (1996) Xerostomia: prevalence and pharmacotherapy. With special reference to beta-adrenoceptor antagonists. *Swedish Dental Journal* **116** (Suppl.), 1–70.
- Nelson, M., Black, A. E., Morris, J. A. & Cole, T. J. (1989) Between- and within-subject variation in nutrient intake from infancy to old age: estimating the number of days required to rank dietary intakes with desired precision. *American Journal of Clinical Nutrition* **50**, 155–167.
- Nishida, N., Tanaka, M., Hayashi, N., Nagata, H., Takeshita, T., Nakayama, K., Morimoto, K. & Shizukuishi, S. (2005) Determination of smoking and obesity as periodontitis risks using the classification and regression tree method. *Journal of Periodontology* **76**, 923–928.
- Noack, B., Jachmann, I., Roscher, S., Sieber, L., Kopprasch, S., Luck, C., Hanefeld, M. & Hoffmann, T. (2000) Metabolic diseases and their possible link to risk indicators of periodontitis. *Journal of Periodontology* **71**, 898–903.
- Norderyd, O. & Hugoson, A. (1998) Risk of severe periodontal disease in a Swedish adult population. A cross-sectional study. *Journal of Clinical Periodontology* **25**, 1022–1028.
- Olaiz, G., Rojas, R., Barquera, S., Shamah, T., Aguilar, C., Cravioto, P., López, P., Hernández, M., Tapia, R. & Sepúlveda, J. (2003) *Encuesta Nacional de Salud 2000. Tomo 2. La salud de los adultos*. Cuernavaca, Morelos: Instituto Nacional de Salud Pública.
- Page, R. C., Krall, E. A., Martin, J., Mancl, L. & Garcia, R. I. (2002) Validity and accuracy of a risk calculator in predicting periodontal disease. *Journal of the American Dental Association* **133**, 569–576.
- Paquette, D. W. (2004) The periodontal-cardiovascular link. *Compendium of Continuing Education in Dentistry* **25**, 681–692.
- Paulander, J., Axelsson, P. & Lindhe, J. (2003) Association between level of education and oral health status in 35-, 50-, 65- and 75-year-olds. *Journal of Clinical Periodontology* **30**, 697–704.
- Paulander, J., Axelsson, P., Lindhe, J. & Wennstrom, J. (2004) Some characteristics of 50/55-year-old individuals with various experience of destructive periodontal disease: a cross-sectional study. *Acta Odontologica Scandinavica* **62**, 199–206.
- Persson, G. R., Persson, R. E., Hollender, L. G. & Kiyak, H. A. (2004) The impact of ethnicity, gender, and marital status on periodontal and systemic health of older subjects in the Trials to Enhance Elders' Teeth and Oral Health (TEETH). *Journal of Periodontology* **75**, 817–823.
- Persson, R. E., Hollender, L. G., MacEntee, M. I., Wyatt, C. C., Kiyak, H. A. & Persson, G. R. (2003) Assessment of periodontal conditions and systemic disease in older subjects. *Journal of Clinical Periodontology* **30**, 207–213.
- Persson, R. E., Kiyak, H. A., Wyatt, C. C., Macentee, M. & Persson, G. R. (2005) Smoking, a weak predictor of periodontitis in older adults. *Journal of Clinical Periodontology* **32**, 512–517.
- Pussinen, P. J., Alftan, G., Rissanen, H., Reunanen, A., Asikainen, S. & Knekt, P. (2004) Antibodies to periodontal pathogens and stroke risk. *Stroke* **35**, 2020–2023.

- Ronderos, M., Pihlstrom, B. L. & Hodges, J. S. (2001) Periodontal disease among indigenous people in the Amazon rain forest. *Journal of Clinical Periodontology* **28**, 995–1003.
- Rozier, R. G. & Beck, J. D. (1991) Epidemiology of oral diseases. *Current Opinion in Dentistry* **1**, 308–315.
- Saito, T., Shimazaki, Y., Koga, T., Tsuzuki, M. & Ohshima, A. (2001) Relationship between upper body obesity and periodontitis. *Journal of Dental Research* **80**, 1631–1636.
- Sbaraglia, M., Turnbull, R. S. & Locker, D. (2002) Risk indicators for periodontal disease in a remote Canadian community – a dental practice-based study. *Journal of Public Health Dentistry* **62**, 51–56.
- Scannapieco, F. A., Bush, R. B. & Paju, S. (2003) Associations between periodontal disease and risk for nosocomial bacterial pneumonia and chronic obstructive pulmonary disease. A systematic review. *Annals of Periodontology* **8**, 54–69.
- Secretaría de Salud (1995) *Encuesta Nacional de enfermedades crónicas 1993*, 2a edición. México, D.F. SSA: Dirección General de Epidemiología.
- Serne, E. H., Gans, R. O., ter Maaten, J. C., Tangelder, G. J., Donker, A. J. & Stehouwer, C. D. (2001) Impaired skin capillary recruitment in essential hypertension is caused by both functional and structural capillary rarefaction. *Hypertension* **38**, 238–242.
- Skrepcinski, F. B. & Niendorff, W. J. (2000) Periodontal disease in American Indians and Alaska Natives. *Journal of Public Health Dentistry* **60** Suppl 1, 261–266.
- Slade, G. D., Ghezzi, E. M., Heiss, G., Beck, J. D., Riche, E. & Offenbacher, S. (2003) Relationship between periodontal disease and C-reactive protein among adults in the atherosclerosis risk in communities study. *Archives of Internal Medicine* **163**, 1172–1179.
- Stern, M. P., Knapp, J. A., Hazuda, H. P., Haffner, S. M., Patterson, J. K. & Mitchell, B. D. (1991) Genetic and environmental determinants of type II diabetes in Mexican Americans. Is there a “descending limb” to the modernization/diabetes relationship? *Diabetes Care* **14**, 649–654.
- Susin, C., Dalla Vecchia, C. F., Oppermann, R. V., Haugejorden, O. & Albandar, J. M. (2004) Periodontal attachment loss in an urban population of Brazilian adults: effect of demographic, behavioral, and environmental risk indicators. *Journal of Periodontology* **75**, 1033–1041.
- Taani, D. S. (2004) Oral health in Jordan. *International Dental Journal* **54**, 395–400.
- Taguchi, A., Sanada, M., Sueti, Y., Ohtsuka, M., Lee, K., Tanimoto, K., Tsuda, M., Ohama, K., Yoshizumi, M. & Higashi, Y. (2004) Tooth loss is associated with an increased risk of hypertension in postmenopausal women. *Hypertension* **43**, 1297–1300.
- Tapia-conyer, R. (1993) *Encuesta Nacional de Enfermedades Crónicas*. México D.F: Secretaría de Salud *Epidemiología*, pp. 25–28.
- Teng, H. C., Lee, C. H., Hung, H. C., Tsai, C. C., Chang, Y. Y., Yang, Y. H., Lu, C. T., Yen, Y. Y. & Wu, Y. M. (2003) Lifestyle and psychosocial factors associated with chronic periodontitis in Taiwanese adults. *Journal of Periodontology* **74**, 1169–1175.
- Thomson, W. M., Slade, G. D., Beck, J. D., Elter, J. R., Spencer, A. J. & Chalmers, J. M. (2004) Incidence of periodontal attachment loss over 5 years among older South Australians. *Journal of Clinical Periodontology* **31**, 119–125.
- Tomita, N. E., Chinellato, L. E., Pernambuco, R. A., Lauris, J. R. & Franco, L. J. (2002) Periodontal conditions and diabetes mellitus in the Japanese-Brazilian population. *Revista de Saude Publica* **36**, 607–613.
- Torrunguang, K., Tamsailom, S., Rojanasomsith, K., Sutthibhaisal, S., Nisapakulorn, K., Vanichjakvong, O., Prapakamol, S., Premisrinirund, T., Pusiri, T., Jaratkulangkoon, O., Unkurapinun, N. & Sritara, P. (2005) Risk indicators of periodontal disease in older Thai adults. *Journal of Periodontology* **76**, 558–565.
- Tsai, C., Hayes, C. & Taylor, G. W. (2002) Glycemic control of type 2 diabetes and severe periodontal disease in the US adult population. *Community Dentistry and Oral Epidemiology* **30**, 182–192.
- Van Staveren, W. A. (1988) *Nutrition and the elderly: A European collaborative study in conjunction with the WHO Special Programme for Research on Aging and International Union of Nutritional Sciences Committee of Geriatric Nutrition. Manual of Operations, Waperningen, The Netherlands. Euronut Report*. Waperningen.
- Varenne, B., Petersen, P. E. & Ouattara, S. (2004) Oral health status of children and adults in urban and rural areas of Burkina Faso, Africa. *International Dental Journal* **54**, 83–89.
- Velázquez, M. O., Rosas, P. M., Lara, E. A. & Pastelín, H. G. (2002) Hipertensión arterial en México: resultados de la Encuesta Nacional de Salud (ENSA) 2000. *Archivos del Instituto de Cardiología de México* **72**, 71–84.
- Willershausen, B., Krahwinkel, T., Valentini, L. & Todt, M. (2003) Correlation between inflammatory periodontal diseases and cardiovascular diseases. *European Journal of Medical Research* **8**, 499–504.
- Wood, N., Johnson, R. B. & Streckfus, C. F. (2003) Comparison of body composition and periodontal disease using nutritional assessment techniques: third National Health and Nutrition Examination Survey (NHANES III). *Journal of Clinical Periodontology* **30**, 321–327.
- World Health Organization (1985) Diabetes mellitus. Report of a WHO Study Group. *World Health Organ Technical Report Series* **727**, 1–113.
- Yoshihara, A., Seida, Y., Hanada, N. & Miyazaki, H. (2004) A longitudinal study of the relationship between periodontal disease and bone mineral density in community-dwelling older adults. *Journal of Clinical Periodontology* **31**, 680–684.
- Zielinski, M. B., Fedele, D., Forman, L. J. & Pomerantz, S. C. (2002) Oral health in the elderly with non-insulin-dependent diabetes mellitus. *Special Care in Dentistry* **22**, 94–98.

Address:

S. Aída Borges-Yáñez
Circuito Exterior S/N
Ciudad Universitaria
Delegación Coyoacán CP 04510
Distrito Federal México
Fax: +52 55 555 03497
E-mail: aborges@servidor.unam.mx

Clinical Relevance

Scientific rationale: The relationship between periodontal health and systemic/general health has been insufficiently characterized, in particular in developing countries.

Principal findings: After controlling for age, sex, general health, and

calculus, urban residence remained a risk marker for periodontitis. As rural communities increasingly replicate the urban lifestyle, and more people become urban dwellers, periodontal deterioration and its relationship with general health may become more apparent.

Practical implications: It is important to improve oral health care using individualized and group-level interventions based on sound evidence, with dental professionals being knowledgeable about coexistence of periodontal and systemic conditions.